Quatromyicide II Disinfectant & Sanitizer



Applications:

Dilute according to the dilution rates mentioned on the label in potable water. Apply using a mop, sponge, cloth or trigger sprayer.

Disinfection & Sanitizing: This product is an effective sanitizer and disinfectant, when used at the dilution rates indicated, on visibly clean, hard, nonporous, inanimate surfaces. As this product's dilution rate and procedure differs according to its application, please refer to the label for the correct usage for this product.

Benefits:

- Disinfects, sanitizes, and deodorizes
- · Effectively removes mold and mildew
- Versatile dilute according to your cleaning application
- · Ideal for the food industry and delicate personnel settings such as daycares or senior care facilities
- Suitable for all washable hard surfaces
- Available as concentrate or ready-to-use

Codes:

Product Code:	52891
• UPC:	0 56069 52891 7

Dilution Rates:

750 ml

5 L

Third Sink Sanitizing 60 ml / 30 L Sanitizing 1:500 Disinfection 1:250 **QUATROMYICIDE II DUSTBANE Available Format: Packaging:** * Refer to Quatromyicide II Bottle: 6.1" x 6.1" x 12.6" (15.5 x 15.5 x 31.9 cm) RTLLSnec Sheet 12.25 lbs (5.56 kg) Box: 12.4" x 12.4" x 13" 49 lbs (22.23 kg) 4 per case

Specifications:

Colourless
02321963
Mild
7.5 - 8.0
Not determined
No
No
Exempt
No
Yes
Yes
Expiry Date

Quatromyicide II Microbiology Efficacy Data Summary



Quatromyicide II kills Pseudomonas aerginosa (Pseudomonas), Staphylococcus aureus (Staph), Salmonella choleraesuis (Salmonella), Campylobacter jejuni (Campylobacter), Listeria monocytogenes (Listeria), Escherichia coli (E. coli), Escherichia coli pathogenic strain 0157:H7 (pathogenic E. coli), Klebsiella pneumoniae (Klebsiella), Pseudomonas cepacia (Pseudomonas), Salmonella typhi (Salmonella), Herpes Simplex Virus 1 & 2 (Herpes), Vaccinia (Pox Virus), Trichophyton mentagrophytes (The Athlete's Foot Fungus).

Quatromyicide II reduces 99.9%Escherichia coli, pathogenic Escheria coli 0157:H7, Listeria monocytogenes, Staphylocuccus aureus and Klebsiella pneumoniae in 60 seconds. At 8 mL per 4 L of water dilution (200 ppm) QUATROMYICIDE II is an effective sanitizer against Salmonella chloeraesuis (Salmonella), Escherichia coli, Shigella sonnei and Staphylococcus aureus, Yersinia enterocolitica, Campylobacter jejuni, Escherichia coli 0157:H7 and Listeria monocytogenes as determined by the AOCA germicide and detergent sanitizer method in the presence of 400 ppm hardness.

Quatromyicide II is an effective fungicide against Trichophyton mentagrophytes (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas, exercise facilities. Quatromyicide II is effective against Avian Influenza virus. Avian Influenza virus. Avian Influenza virus, Pseudorabies virus, Canine distemper virus, Newcastles Disease, Marek's Disease Virus, Infectious Bronchitis Arkansas 99'Virus, Human Immunodeficiency Virus (HIV), Adenovirus Type 5, and Infectious Laryngotracheitis Virus.

Quatromyicide II is a Hospital Disinfectant at 18 mL per 4 L (450 ppm active), Bacteriacidal according to the current AOAC Disinfectants Use-Dilution Method, Fungicidal according to the AOAC fungicidal test and Virucidal* according to the virucidal qualification, modified in the presence of 5% organic serum against: Campylobacter jejuni, Pseudomonas aeruginosa, Staphylococcus aureus, Salmonella choleraesuis, Salmonella typhi, Escherichia coli pathogenic strain 0157:H7, Trichophyton, mentagrophytes, Yerisinia entercolitica, *Influenza A2/Hong Kong (H1N1), Vaccinia, Human Immunodeficiency Virus, Adenovirus Type 5, Infectious Laryngotracheitis Virus, Klebsiella pneumoniae, Listeria monocytogenes, Pseudomonas cepacia, Escherichia coli, Herpes Simplex Type 1, Herpes Simplex Type 2